

Biologia Geral e Experimental

Universidade Federal de Sergipe

Biol. Geral Exper., São Cristóvão, SE 7(1):19-28

13.ix.2007

NOTES ON THE REPRODUCTION, BEHAVIOUR AND DIET OF *SAGUINUS NIGER* (PRIMATES: CALLITRICHIDAE) IN A FOREST REMNANT AT THE NATIONAL PRIMATE CENTRE, ANANINDEUA, PARÁ

Suleima do Socorro Bastos da Silva¹

Stephen Francis Ferrari²

ABSTRACT

This study presents data on the reproduction, behaviour and diet of a free-ranging group of black-handed tamarins, *Saguinus niger*, in an urbanised forest at the National Primate Centre in Ananindeua, eastern Brazilian Amazonia. The study group contained between five and seven members, changing due to the disappearance of an adult and the births of three infants. Quantitative behavioural data were collected in scan samples during five days each month between May and September 2000, covering the transition between wet and dry seasons. The general activity budget (n = 935 records) was 58.7% locomotion, 16.2% feed, 10.1% rest, 9.7% forage, and 5.3% social behaviour/others. During foraging, subjects spent 83.9% of their time scanning visually for signs of prey, and only 16.1% manipulating substrates. The diet was frugivorous-insectivorous, with 94.2% of feeding records involving fruit, and 5.8% insects. Nine of the plant species were identified, four of which were previously unrecorded for *S. niger*. All the identified insect prey were orthopterans. The group spent most of its time in the middle to lower forest strata moving on substrates of small diameter. Despite the overall limitations of the data collected, then, the study group exhibited typical tamarin patterns of behaviour, and was clearly well adapted to conditions at the study site, despite the degree of habitat disturbance.

Key words: *Saguinus niger*, behaviour, ecology, diet, habitat fragmentation, Amazonia.

RESUMO

Este estudo apresenta dados sobre a reprodução, o comportamento e a dieta de um grupo silvestre de *Saguinus niger*, em um fragmento florestal urbano no Centro Nacional de Primatas em Ananindeua, Pará (Amazônia oriental). O grupo de estudo variou entre cinco e sete membros, mudando pelo desaparecimento de um adulto e o nascimento de três infantes. Dados quantitativos de comportamento foram coletados através de amostragem de varredura durante cinco dias por mês, entre maio e setembro de 2000, abrangendo a transição entre as estações chuvosa e seca. O orçamento geral de atividades (n = 935 registros) foi de 58,7% de locomoção, 16,2% de alimentação, 10,1% de descanso, 9,7% de forrageio e 5,3% comportamento social/outros. Durante o forrageio, os indivíduos gastaram 83,9% de seu tempo procurando visualmente por presas, e somente 16,1% manipulando substratos. A dieta foi frugívora-insetívora, com 94,2% de registros alimentares envolvendo frutos e 5,8% insetos. Nove espécies de plantas foram identificadas, das quais quatro não tinham sido registradas para *S. niger*. Todos os insetos consumidos foram da ordem Orthoptera. O grupo passou a maior parte de seu tempo nos estratos florestais medianos, utilizando substratos finos. Apesar de todas as limitações dos dados coletados, o grupo de estudo exibiu padrões comportamentais típicos do gênero, e está claramente bem adaptado às condições da área de estudo, apesar do grau de perturbação do habitat.

Palavras-chave: *Saguinus niger*, comportamento, ecologia, dieta, fragmentação de habitat, Amazônia.

¹ Centro de Ciências Biológicas, Universidade Federal do Pará, Rua Augusto Correa 1 - Campus Universitário Guamá, CEP 66075-110, Belém, Pa, suleima_silva@yahoo.com.br.

² Universidade Federal de Sergipe, Av. Marechal Rondon s/n, São Cristóvão, Jardim Rosa Elze, CEP 49100-000, ferrari@pq.cnpq.br.

INTRODUCTION

Remnant populations of primates isolated in fragments of habitat are relatively vulnerable to catastrophic events such as fires or epidemics, and may suffer inbreeding depression, all of which increase considerably the risks of local extinction (Lande & Barrowclough, 1987; Lacy, 1997). Such populations nevertheless play an increasingly important role in the conservation of many species, especially in the Atlantic Forest (Ferrari & Diego, 1995; Ruiz-Miranda et al., 2002; Jerusalinsky et al., 2006), but also in Amazonia (Silva, 2003; Gordo, 2005). However, the development of effective management strategies depends on the availability of reliable data on the ecology of the species, primarily in the context of habitat fragmentation.

The callitrichids (marmosets and tamarins) are among the most ecologically tolerant platyrrhines, and many species exhibit a clear preference for disturbed habitat in comparison with primary forest. Some species, such as the common marmoset (*Callithrix jacchus*) may even be abundant in urban environments (Cunha et al., 2006). Morphological specialisations for this “colonising” niche include small body size (Martin, 1992; Ferrari, 1993), which contributes to reduced spatial needs in comparison with larger-bodied primates. The success of callitrichids in marginal habitats may also be reinforced by the lack or absence of competition from these larger, ecologically less tolerant platyrrhines.

Despite being distributed throughout much of southeastern Amazonia (Hirsch *et al.*, 2002), the ecology of *Saguinus niger* is still poorly known, although the available evidence (Oliveira, 1996; Oliveira & Ferrari, 2000; Veracini, 2000) indicates that the behavioural ecology of this tamarin is typical of the genus, and that the species is essentially frugivorous-insectivorous. Data on both *S. niger* (Ferrari & Lopes, 1990, 1996; Veracini, 2000) and the

closely-related *Saguinus midas* (Egler, 1993) indicate that these species may be ecologically more successful than sympatric callitrichids, *Mico argentatus* and *Saguinus bicolor*, respectively. This implies that they may also be more tolerant of habitat disturbance, and in fact Egler (1993) concluded that habitat fragmentation east of Manaus has favoured expansion of the geographic range of *S. midas* into areas previously occupied by *S. bicolor*. Obviously, this process has deleterious implications for the conservation of *S. bicolor* – one of the most endangered of Amazonian primates – and there is an urgent need for more data on all aspects of the ecology of the two species (Gordo, 2005).

With this in mind, this study presents preliminary data on the behavior and diet of a free-ranging group of black-handed tamarins (*S. niger*) in an isolated forest fragment located within an urban environment in eastern Pará. In addition to providing new data on this poorly-known platyrrhine, it is hoped that the study will contribute to the development of effective management strategies for fragmented populations of tamarins throughout the Amazon.

METHODS

Study site: The present study took place in an isolated fragment of approximately 25 hectares of Amazonian *terra firme* forest on the grounds of the National Primate Centre (CENP) in Ananindeua, Pará (01°21' S, 48°22' W). The forest has been logged selectively, but can be characterised as disturbed primary habitat (O. Cardoso, pers. comm.), with trees of up to 30 m and clearings dominated by palms and epiphytes. The area is criss-crossed by a system of trails, some of which are paved. The site is inhabited by one other primate (*Saimiri sciureus*), and a number of small- to medium-sized mammals, such as pacas (*Agouti paca*), sloths (*Bradypus tridactylus*), and squirrels (*Sciurus* sp.).

Data collection: A preliminary study took place between January and April, 2000, period that coincides with the first half of the wet season in the study area (monthly precipitation peaked at 481 mm in April). During this period, study group members were identified and habituated to the presence of human observers, and data collection procedures were tested. As the group was small in size, it was possible to distinguish members on the basis of body size and sex.

Between May and September (the dry season normally starts in July-August, although precipitation was never less 100 mm in any given month), quantitative behavioural data were collected in instantaneous scan samples (Martin & Bateson, 1993), using the sampling schedule recommended for callitrichids by Ferrari & Rylands (1994), and also used by Oliveira (1996) and Veracini (2000) in their field studies of *S. niger*. This schedule involves the collection of a one-minute sample at intervals of five minutes. Data were collected on five days each month, with observation sessions lasting between four and seven hours.

At the beginning of each scan sample, the time and location was recorded and, for each subject observed during the scan, the following information was collected: its activity state (categories: travel, forage, feed, rest, social behaviour, and miscellaneous); identity or age-sex class; its height above the ground (m) and the diameter (cm) of the supporting substrate; any other information considered relevant. The foraging category was divided into two subcategories: visual scanning for signs of prey, and manipulation of substrates. The set of information collected for a given subject during a scan is referred to here as a record. Complementary behavioural data were collected throughout the study period, in *ad libitum* (Altmann, 1974) fashion.

The group's activity budget was estimated according to the relative proportions of records of different behaviour categories. The composition of the

diet was based on the relative proportions of feeding records attributed to different food items. Whenever subjects were observed ingesting fruit, the source was marked, and samples were collected for taxonomic identification. Defecated seeds were also collected for identification and measurement. Insects were identified, whenever possible, through the collection of prey parts discarded by the subjects. For the analysis of vertical stratification, records of subject height were grouped in classes of 5 m, following Oliveira (1996).

RESULTS

Group composition: At the beginning of the preliminary study, the group had five members, including three adults (two males and one female), a subadult female and a juvenile male. This increased to six with the birth of a single female infant in early January (no later than the 13th). In June, one of the adult males was captured by a resident of the housing estate located next to the CENP. This animal was released back into the CENP on request of the first author, but it subsequently disappeared, leaving the study group with a total of five members (Table 1).

By the end of July (no later than the 24th), a pair of twins were born, increasing group size to seven members. The infants were first observed on the back of the mother, which was having some difficulty moving and, at one point, was left hanging by her hands and feet, with her back – and the infants – facing the ground. Subsequently, other group members – in particular the adult male – carried infants, a typical behaviour pattern (Digby *et al.*, 2007), although normally only one at a time. During approximately two-thirds of the time, at least one of the infants remained with the mother. Transfers of infants were invariably stimulated by the prospective carrier, by approaching the infant.

By September, these twins were already exploring the environment independently of adult caregivers, principally during feeding bouts. Typically, during feeding bouts, the infants would explore the tree crown on their own, but would be retrieved by one of the adults when the group moved on. During the daily “siesta” period, around midday, the infants engaged in play behaviour, including the attempted capture of butterflies, and chasing small birds.

These infants begged food items from the adults and subadult in typical callitrichid fashion (Tardif et al., 1993), with shrill and insistent vocalisations, and invariably succeeded in obtaining the transfer of the desired food item. While this behaviour pattern is typical of all callitrichids, Oliveira (1996) only recorded food transfer between the mother and her infant in a group of *S. niger*, although this may have been because the infant had only just started to move independently at the time of monitoring.

Activity budget Monitoring of the study group was affected by a number of problems, including the capture of the adult male in June, which resulted in group members being more wary of observers during the following days. In subsequent months, which also coincided with major building work within the compound, the group spent increasingly more time – often more than four hours – in a sector of the fragment covered in extremely dense vegetation which had no access trails, as well as visiting the back gardens of houses that adjoin the 5-metre high perimeter wall of the CENP compound.

If monitoring was impaired in the first case, it became impossible in the second, given that there was no way of accompanying the animals beyond the perimeter wall. It wasn’t clear what resources were exploited in the gardens, but occasional shouts heard from neighbours indicate that they may even have entered some of the houses in search of food, a behaviour observed in some other urban callitrichids (SFF, pers. obs.). These visits lasted from 30 to 75

minutes, and were often continuous with time spent in the denser forest. In addition to these problems, administrative restrictions at the site limited data collection to periods of no more than seven hours, rather than continuous monitoring throughout the daily activity period, as originally planned.

Despite all these problems, quantitative data were recorded in a total of 424 scan samples, during which 935 records were collected. While the average number of records collected per scan (2.2) may be considered adequate, considering the size of the study group (Ferrari & Rylands, 1994), the small overall number of records limits conclusions on behaviour patterns. Despite these obvious limitations, the activity budget was typical of tamarin species, with the vast majority of activity time being devoted to locomotion, foraging and feeding (Figure 1). However, it seems likely that locomotion is overestimated relative to the other categories, considering both the observation conditions, as described above, and the intrinsic bias of the scan sampling procedure (Ferrari & Rylands, 1994).

During foraging ($n = 150$ records), 83.9% of time was spent scanning visually for signs of prey, the remainder being dedicated to the manipulation of substrates. Generally, live leaves and twigs were handled superficially, whereas dead leaves were opened destructively. Holes in the trunks of trees and the forks of branches were also investigated intensively.

Social behaviour: Four categories of social behaviour were recorded during the present study: allogrooming, agonism, play, and scent marking. Scent marking was the most frequently observed category, and almost all observations were of sternal and suprapubic marking. Typically, two or three individuals – normally adults – would mark the same place on the substrate consecutively. Circumgenital marking was rare ($n = 3$ records, two of which by juveniles), and did not involve urination.

Play was also a common behaviour, and invariably involved the immature individuals, which engaged in tag-chasing. On three occasions, birds (two small passeriformes and an immature toucan, *Ramphastos* sp.) were involved in the play, with the tamarins hiding behind branches and then jumping towards the birds. The passeriformes responded by hiding behind branches, whereas the toucan flew away immediately.

Allogrooming was observed primarily during rest periods, during the middle of the day, but were normally very discrete, reducing even further the visibility of the subjects. Juveniles generally groomed adults more than vice versa. In one case, the subadult female groomed the group's oldest adult male, who then inspected and sniffed her genitalia, while she adopted a solicit-grooming posture. At this moment, the male bit the female lightly on the ear and moved away. Agonism was restricted to occasional exchanges of excited vocalisations.

Diet: Once again, the relatively small number of feeding records collected ($n = 85$) limits the analysis of the study group's diet. Despite this, the subjects were clearly frugivorous-insectivorous, dedicating 94.2% of feeding time to the ingestion of fruit, and 5.8% to the predation of insects. While plant exudates (gum) are a dietary staple for many callitrichids, the lack of consumption in the present study may be due to the period of the present study, given that in the previous study, gum was only exploited at end of the dry season (November and December), when fruit was scarce.

A total of nine tree species were exploited for their fruit (Table 2), only four of which had been recorded previously (Oliveira & Ferrari, 2000; Veracini, 2000). In four cases, the fruit pulp was ingested together with the seeds, which were defecated intact. It is interesting to note here that the largest seed ingested by *S. niger* in the study of Oliveira & Ferrari (2000) were 18 mm in length, whereas many seeds of larger size (more than 20 mm in length) were not

ingested. Only five insect prey were identified, and all belonged to the order Orthoptera, the archetypal callitrichid prey. Of these, three were grasshoppers (Acrididae), and two were crickets (Grillidae).

Vertical Spacing: Study group members were observed at all different levels within in the forest, but clearly preferred the middle to lower strata (Figure 2). Similar preferences have been recorded at other sites (Lopes, 1993; Oliveira, 1996; Bobadilla & Ferrari, 2000; Veracini, 2000). Despite being active at the lower levels in the forest, the subjects came to the ground only very rarely, and only to cross gaps in the forest.

This is consistent with typical tamarin behaviour, and also the preference of larger-bodied sympatric primates, such as *Alouatta* and *Chiropotes* for the higher strata (Bobadilla & Ferrari, 2000). Veracini (2000) recorded a very similar pattern, including the relative predominance of foraging behaviour in the 5-15 m stratum. Substrate sizes (Figure 3) reflect the preference for relatively small branches in the middle strata, and the relatively infrequent practice of the vertical clinging and leaping mode of locomotion on large, vertical trunks.

DISCUSSION

Behaviour patterns are a fundamental component of the ecology of any primate species, and vary systematically in accordance with local conditions and seasonal fluctuations in resources. Understanding basic patterns and the response to shifts in resource availability provides important insights into evolutionary processes and ecological specialisations. The tamarins (*Saguinus* spp.), essentially Amazonian in their distribution (Hirsch *et al.*, 2002), are highly active frugivore-insectivores, and spend a large proportion of their time foraging and moving rapidly through the forest.

Comparisons among studies are hampered by differences in variables such as sampling procedures, study duration and timing, and group composition (Ferrari & Rylands, 1994). In the case of *S. niger*, all three studies conducted to date have followed the same sampling schedule recommended by these authors, and in this respect, are directly comparable (Table 3). While there are broad overall similarities, there are also notable differences, such as relatively high rates of locomotion in the CENP group, and low levels of social behaviour in Veracini's ECFPn group. Increased locomotion in the CENP group almost certainly reflects observation conditions, as indicated by differences in the number of records collected (for similar periods of data collection). By contrast, the time spent feeding was remarkably consistent across studies, which may reflect the relative importance of this activity.

While diets were also broadly similar, with regard to the predominance of fruit and the relatively minor contribution of arthropods (Table 4), Veracini's (2000) data are divergent in terms of the variety of plant resources exploited. A number of different factors may have contributed to this difference, including forest type, season, and in particular, sympatry with *M. argentatus*. The latter species occurs only at ECFPn, where it feeds intensively on exudates by gouging the bark of gum-producing species such as *Parkia ulei* (Mimosaceae). The behaviour of the marmosets thus provides an alternative resource for *S. niger*, which may be especially important during the dry season. The site at ECFPn was also within an area of continuous forest (rather than a fragment as at the other sites), with extensive river edge habitat and high concentrations of *Symphonia globulifera* (Guttiferae), a major source of nectar for many callitrichids. Most tamarins will consume exudates when or where they are available (Digby *et al.*, 2007), and sympatry with marmosets may be a significant factor in the ecology of a tamarin species (Lopes & Ferrari, 1994).

Despite its methodological limitations, then, the present study has shown that *S. niger* is not only able to survive and reproduce in a highly-disturbed forest matrix, it may do this without major alterations in behaviour patterns. This is promising for the long-term survival of the species, and eventual management of remnant populations in a critically impacted region.

Acknowledgements: We are grateful to the staff of the National Primate Centre, in particular its director, José Augusto Muniz, for their support during data collection. SFF acknowledges the Brazilian Scientific Research Council (CNPq) for a research fellowship (Process no. 307506/2003-7).

REFERENCES

- Altmann, J. 1974. Observational study of behaviour: sampling methods. **Behaviour** 49:227-267.
- Bobadilla, U.L. & S.F. Ferrari, 2000. Habitat use by *Chiropotes satanás utahicki* and syntopic platyrrhines in eastern Amazonia. **American Journal of Primatology** 50(3):215-224.
- Cunha, A.A., M.V. Vieira & C.E.V. Grelle, 2006. Preliminary observations on habitat, support use and diet in two non-native primates in an urban Atlantic forest fragment: the capuchin monkey (*Cebus* sp.) and the common marmoset (*Callithrix jacchus*) in the Tijuca forest, Rio de Janeiro. **Urban Ecosystems** 9(4):351-359.
- Digby, L.J., S.F. Ferrari & W. Saltzmann, 2007. Callitrichines: the role of competition in cooperatively breeding species, pp. 85-106. *In: Primates in Perspective* (C.J. Campbell, A. Fuentes, K.C. MacKinnon, M. Panger & S.K. Bearder, Eds.). Oxford University Press.
- Egler, S.G. 1993. First field study of the pied tamarin, *Saguinus bicolor bicolor*. **Neotropical Primates** 1(2):13-14.
- Ferrari, S.F. 1993. Ecological differentiation in the Callitrichidae, pp. 314-328. *In: Marmosets and tamarins: systematics, behaviour, and ecology* (A.B. Rylands, Ed.). Oxford University Press.
- Ferrari, S.F. & V.H. Diego, 1995. Habitat fragmentation and primate conservation in the Atlantic Forest of eastern Minas Gerais, Brazil. **Oryx** 29(3):192-196.
- Ferrari, S.F. & A.B. Rylands, 1994. Activity budgets and differential visibility in field studies of three marmosets (*Callithrix* spp.). **Folia Primatologica** 63(2):78-83.
- Ferrari, S.F. & M.A. Lopes, 1990. A survey of primates in central Pará. **Boletim do Museu Paraense Emílio Goeldi, Série Zoologia**, 6(2):169-179.

- Ferrari, S.F. & M.A. Lopes, 1996. Primates populations in eastern Amazonia, pp. 53-68. *In: Adaptive Radiations of Neotropical Primates* (M.A. Norconk, A.L. Rosenberger & P.A. Garber, Eds.). Plenum Press.
- Gordo, M. 2005. A conservação de *Saguinus bicolor* em fragmentos florestais na cidade de Manaus, Amazonas. **Livro de Resumos do XI Congresso Brasileiro de Primatologia**, p. 50.
- Hirsch, A., L.G. Dias, L.O. Martins, R.F. Campos, E.C. Landau & N.A.T. Resende, 2002. BDGEOPRIM – Database of geo-referenced localities of Neotropical primates. **Neotropical Primates** 10(2):79-84.
- Jerusalinsky, L., M.M. Oliveira, R.F. Pereira, V. Santana, P.C. Bastos & S.F. Ferrari, 2006. Preliminary evaluation of the conservation status of *Callicebus coimbrai* Kobayashi & Langguth, 1999, in the Brazilian State of Sergipe. **Primate Conservation** 21:25-32.
- Lacy, R.C. 1997. Importance of genetic variation to the variability of mammalian populations. **Journal of Mammalogy** 78:320-335.
- Lande, R. & G. Barrowclough, 1987. Effective population size, genetic variation, and their use in population management, pp. 87-123. *In: Viable populations for conservation* (M.E. Soulé, Ed.). Cambridge University Press.
- Lopes, M.A. 1993. Conservação do cuxiú-preto, *Chiropotes satanas satanas* (Platyrrhini, Primates) e de outros mamíferos na amazônia oriental. **Dissertação de Mestrado**, Centro de Ciências Biológicas, Universidade Federal do Pará.
- Lopes, M.A. & S.F. Ferrari, 1994. Foraging behaviour of a tamarin group (*Saguinus fuscicollis weddelli*), and interactions with marmosets (*Callithrix emiliae*). **International Journal of Primatology** 15:373-387.
- Martin, R.D. 1992. Goeldi and the dwarfs: the evolutionary biology of the small New World primates. **Journal of Human Evolution** 22:367-393.
- Martin, P. & P. Bateson, 1993. **Measuring behaviour: an introductory guide**. Cambridge University Press.
- Oliveira, A.C.M. 1996. Ecologia e comportamento alimentar, padrões de atividade e uso de espaço de um grupo de *Saguinus midas niger* (Callitrichidae, Primates) na amazônia oriental. **Dissertação de Mestrado**, Departamento de Psicologia Experimental, Universidade Federal do Pará.
- Oliveira, A.C.M. & S.F. Ferrari, 2000. Seed dispersal by black-handed tamarins, *Saguinus midas niger* (Callitrichidae, Primates): implications for the regeneration of degraded forests in eastern Amazonia. **Journal of Tropical Ecology** 16:709-716.
- Ruiz-Miranda, C.R. 2002. Conservação do mico leão dourado: uma estratégia multidisciplinar, multi-institucional e internacional a longo prazo. **Livro de Resumos do X Congresso Brasileiro de Primatologia**, p. 54.
- Silva, S.S.B. 2003. Comportamento alimentar de cuxiú-preto (*Chiropotes satanas*) na área de influência do reservatório da Usina Hidrelétrica de Tucuruí-PA. **Dissertação de Mestrado**, Departamento de Zoologia, Museu Paraense Emílio Goeldi/Universidade Federal do Pará.
- Tardif, S.D., M.L. Harrison & M.A. Simek, 1993. Communal infant care in marmosets and tamarins: relation to energetics, ecology, and social organization, pp. 220-234. *In: Marmosets and tamarins: systematics, behaviour, and ecology* (A.B. Rylands, Ed.). Oxford University Press.
- Veracini, C. 2000. Dados preliminares sobre a ecologia de *Saguinus niger* na Estação Científica Ferreira Penna, Caxiuanã, Pará, Brasil. **Neotropical Primates** 8(3):108-113.

Table 1. Composition of the study group during the monitoring period.

Month	Adult males	Adult females	Subadult (M/F)	Juvenile (M/F)	Infant
December 1999	2	1	1F	1M	0
January 2000	2	1	1F	1M	1F
June	1	1	1F	1M	1F
July	1	2	1M	1F	2
September	2	2	0	1F	2

M: male; F: female.

Table 2. Tree species exploited for their fruit by the *S. niger* study group at CENP.

Taxon		Seed ingested	Maximum length x width of seed (mm)	Recorded for <i>S.m. niger</i> by
Boraginaceae	<i>Cordia</i> sp.	No		Veracini (2000)
Elaeocarpaceae	<i>Sloanea</i> sp.	Yes	12 x 10	-
Apocynaceae	<i>Lacmellea</i> sp.	No		Veracini (2000)
Chrysobalanaceae	<i>Licania alba</i>	No		-
Ochnaceae	<i>Ouratea</i> sp.	Yes	11 x 8	-
Mimosaceae	<i>Parkia nitida</i>	Yes	23 x 10	-
Moraceae	<i>Pouroma guianensis</i>	No		Oliveira & Ferrari (2000)
Sapotaceae	<i>Pouteria</i> sp.	Yes	14 x 7	Oliveira & Ferrari (2000)
Anacardiaceae	<i>Tapirira guianensis</i>	No		Veracini (2000) ¹

¹Exudate exploited.Table 3. Activity budgets recorded for *S. niger* (all studies followed the same scan sampling schedule).

Study site	% of time spent:					N records	Source
	Travel	Forage	Feed	Rest	Misc.		
CENP	58.70	9.70	16.21	10.06	5.33	935	Present study
Fazenda Vitória	46.08	15.19	16.73	17.46	4.54	3659	Oliveira (1996)
ECFPn	42.38	18.86	17.64	14.50	0.53	1904	Veracini (2000)

Table 4. Composition of the diet of *S. niger* at different study sites.

Study site	% of feeding records:				Source
	RPPs ¹	Exudate	Animal prey	Nectar	
CENP	94.2	-	5.8	-	Present study
Fazenda Vitória	87.5	3.1	9.4	-	Oliveira & Ferrari (2000)
ECFPn	48.8	23.8	4.5	22.9	Veracini (2000)

¹Reproductive Plant Parts (fruit, flowers).




Figure 1. General activity budget of the *S. niger* study group (n = 935 records). The category “other” refers to defecation and urination.




Figura 2. Use of different forest strata by the *S. niger* study group, according to scan sample records (n = 935). “Others” includes social behaviour and defecation/urination.




Figura 3. Use of substrates of different sizes by the *S. niger* study group, according to scan sample records (n=935).