

Biologia Geral e Experimental

Universidade Federal de Sergipe

Biol. Geral Exper., São Cristóvão, SE 7(2):33-40

30.xii.2007

EFFECTS OF THE AQUEOUS EXTRACTS OF PLANTS OF THE GENERA *APODANTHERA* (CUCURBITACEAE) AND *JATROPHA* (EUPHORBIACEAE) ON THE LETHALITY OF THE VENOM OF *BOTHRUPS JARARACA* (SERPENTES, VIPERIDAE)

Jeane Carvalho Vilar¹

Celso Morato de Carvalho²

Maria de Fátima Domingues Furtado³

ABSTRACT

Four plants popularly used as antiophidic in the cerrado and caatinga regions of the Northeastern and Central Brasil were analyzed in order to verify the inhibition of the venom lethal action of the Neotropical viper *Bothrops jararaca* (Serpentes, Viperidae). The plants were *Apodanthera villosa* (Cucurbitaceae), *Apodanthera glaziovii* (Cucurbitaceae), *Jatropha molissima* (Euphorbiaceae) and *Jatropha elliptica* (Euphorbiaceae). The LD₅₀ of the venom was 37.1 µg/mouse. The control groups were mouse inoculated with the viper venom (71.2 µg), the experimental groups were mouse inoculated with the viper venom (71.2 µg) and aqueous extract of the plants. The extract of *A. villosa* (1.0 mg) increased the survivor time of the experimental animals compared to control groups.

Keywords: Antiophidic plants, *Apodanthera*, *Jatropha*, caatinga, cerrado.

RESUMO

Foram analisadas quatro plantas popularmente utilizadas como antiofídicas nas regiões de caatinga e cerrado do nordeste e Brasil central, para verificar a inibição da ação letal do veneno da serpente *Bothrops jararaca* (Serpentes, Viperidae). As plantas foram *Apodanthera villosa* (Cucurbitaceae), *Apodanthera glaziovii* (Cucurbitaceae), *Jatropha molissima* (Euphorbiaceae) e *Jatropha elliptica* (Euphorbiaceae). A DL₅₀ do veneno foi 37.1 µg/camundongo. O grupo controle foi camundongos inoculados com o veneno de jararaca (71.2 µg), o grupo experimental foi camundongos inoculados com o veneno de jararaca (71.2 µg) e os extratos aquosos das plantas. O extrato de *A. villosa* (1.0 mg) aumentou o tempo de sobrevivência dos animais experimentais quando comparados com o grupo controle.

Palavras-chave: Plantas antiofídicas, *Apodanthera*, *Jatropha*, caatinga, cerrado.

INTRODUCTION

The venom of the snakes – the Neotropical vipers – of the genus *Bothrops* (family Viperidae) has a systemic hemorrhagic effect and increases the coagulation time of the blood (Furtado *et al.*, 1991; Gutierrez, 2002). *Bothrops* are responsible for the majority of snake envenomations that occur in Brasil.

Treatment with antivenins is effective and recommended, but in many remote regions, where medical services are precarious, envenomation is often treated with local plant-based remedies (Otero *et al.*, 2000; Vilar *et al.*, 2004, 2005). Two of the plants recommended for the treatment of envenomation by vipers are known locally as “batata-de-teiú” and “pinhão-bravo” (Braga, 1960; Mors, 1991; Martz,

¹Faculdade Pio Décimo, Aracaju, Sergipe, jcvilar@bol.com.br.

²Instituto Nacional de Pesquisas da Amazônia, Coordenação de Pesquisas em Biologia Aquática, Manaus, Amazonas, cmorato@inpa.gov.br.

³Instituto Butantan, Departamento de Herpetologia, São Paulo, SP, fatiafurtado@butantan.gov.br.

1992), although few reliable reports of the antivenin properties of plants with these names are available (Amuí *et al.*, 2003). A third plant, referred to as “cabeça-de-negro” is also frequently cited as an inhibitor of viper venom, although the report (Nakagawa *et al.*, 1982) does not identify the species, and there are at least ten different plant species known by this name (Silva *et al.*, 1998). Given these reports, the objective of the present study was to contribute to the scientific knowledge and therapeutic use of these plants, in particular with regard to their effectiveness against viper venom.

METHODS

Venom: Mixture of lyophilized *Bothrops jararaca* venom provided by the Instituto Butantan, São Paulo State Health Department. The mixture was stored at -20°C until used.

Animal subjects: Experiments were conducted on non-isogenic Swiss mice weighing 18-22 g supplied by the Central Breeding Laboratory of the Universidade Federal de Sergipe in São Cristóvão.

Plant samples: Four plant species belonging to the Cucurbitaceae and Euphorbiaceae families were analyzed in the present study. The Cucurbitaceae was represented by the herbaceous *Apodanthera villosa* C. Jeffrey, known locally as “batata-de-teiú”, and the climber *Apodanthera glaziovii* Glaziov ex Cogniaux, known as “cabeça-de-negro”. These two species, together with the bush known as “pinhão-bravo” (*Jatropha mollissima* (Pohl) Baill), a member of the Euphorbiaceae, were collected at Curituba (09°41'S, 37°53'W), in the northeastern Brazilian state of Sergipe. A second species of Euphorbiaceae, a herbaceous plant also known as “batata-de-teiú” (*Jatropha elliptica* (Pohl) Muell. Arg.), was collected in Peixe (12°01'S, 48°32'W), in the central Brazilian

state of Tocantins. Specimens were identified at the Department of Botany of the Universidade Federal de Goiás in Goiânia, and the Instituto de Botânica of the São Paulo State Environment Department.

Preparation of the aqueous extract: Tubers of *A. villosa*, *A. glaziovii* and *J. elliptica*, and stems of *J. mollissima* were oven-dried at 40°C, and ground to dust in a mortar. Raw extracts were prepared via the decoction of 200 grams of the durst for 10 minutes. This extract was filtered and lyophilized.

Lethal dose 50% of the venom: The LD₅₀ of the venom was determined by a regression analysis of the number of mice surviving or dying during the 48 hours following inoculation with *Bothrops* venom. The coefficient of regression (y) is given by $y = a + bx$, where x is the logarithm of the dose concentration (Fisher & Yates, 1953). The value of x for 50% mortality is found where $y = 5$. The doses of viper venom were 23.0, 27.6, 33.1, 39.7 and 47.6 µg. The between-dose interval was established based on the LD₅₀ of the venom of *B. jararaca* determined by the Instituto Butantan. Starting with the smallest dose of 23.0µg, which is smaller than the venom's LD₅₀, larger doses were determined by a 1.2 dilution factor, the standard ratio for venom studies. For each dosage, six mice were inoculated intraperitoneally with 0.5 ml of the venom dissolved in 0.9% NaCl solution.

Lethal dose 50% of the plant extracts: It was necessary to determine the LD₅₀ of the plants in order to establish extract dosages for the tests on the inhibition of the lethality of the venom. Three experiments were conducted on the extracts of each of the species *A. villosa*, *A. glaziovii* and *J. mollissima*. The LD₅₀ values of these plants were obtained from the combined analysis of all experiments, derived from the probits. Extract doses varied from 0.004 to 0.008 g, 0.001 to 0.011 g and 0.0003 to 0.04 g, respectively. Analysis of the 50% lethal dose of *J. elliptica* was

based on a single experiment, with doses varying from 0.01 to 0.03 g

Experiments on the inhibition of lethality: The venom's LD_{50} was doubled to determine the challenge dose (Gutiérrez *et al.*, 1990) used as the criterion for testing the ability of the plants to neutralize the toxin. Lethality was tested in two groups of six mice. In the control group, the mice were inoculated with $2LD_{50}$ of the venom dissolved in 0.9% NaCl solution. In the experimental group, the mice received the same challenge dose as the control group, but with the addition of the plant extracts dissolved in 0.9% NaCl. The solutions of venom plus extract were incubated at 37°C for 30 minutes prior to inoculation. The doses for each extract were: *A. villosa* 1.00 mg (LD_{50} 0.018 g), *A. glaziovii* 1.00 - 1.48 - 3.0 - 5.0 mg (LD_{50} 0.007 g), *J. mollissima* 1.0 mg (LD_{50} 0.002 g), and *J. elliptica* 0.74 - 1.00 - 1.48 - 5.0 - 10.0 mg (LD_{50} 0.018 g). These doses were determined after establishing the corresponding LD_{50} values, which were done through probit analysis. Each mouse was inoculated intraperitoneally with 0.5 ml of the solution. To determine survival time, the animals were observed at one-hour intervals during 48 hours.

Data analyses: All probit values were obtained using the method of Fisher (1949) and Fisher & Yates (1953). In the inhibition of lethality experiments, the first step was to confirm the homogeneity of controls and experimental samples for each dosage of the extract. When considered homogeneous, samples were grouped for final analysis, using one-factor ANOVA and the *t* test for comparisons between the means of two samples (Vanzolini, 1993; Zar, 1996).

RESULTS

A total of six control experiments were carried out, whereas the number of extract tests varied

according to the plant species. Regressions and 50% lethal doses were determined for controls and extracts (Table 1). In the case of the control, the regression of lethality on venom dosage was $y = -7.99 + 8.28x$, with a confidence interval of 34.43 ± 40.45 (Table 1). The 50% mortality threshold (LD_{50}) is found at $y = 5$, so in this experiment, the dose of *B. jararaca* venom for 50% mortality of the inoculated mice was estimated as $x = 37.09$ mg. Given this, the challenge dose ($2LD_{50}$) used as the criterion for testing the effects of the extracts on the lethality of the venom was 74.18 mg.

Four experiments were conducted using the extract (1.0 mg) of *A. villosa*. As both control and experimental groups were homogeneous ($F_{0.05(1)5,27} = 1.436$, $p > 0.05$ and $F_{0.05(1)3,42} = 0.857$, $p > 0.05$, Table 2), samples were grouped and compared using the *t* test. Survival time was significantly greater in the experimental group ($t_{0.05(1),67} = 2.084$, $p < 0.05$, Table 3), which indicates that the extract of this species was effective in inhibiting the lethality of the viper venom.

The extract (1.0 mg) of *A. glaziovii* was tested in four experiments, the results of which were homogeneous among samples ($F_{0.05(1)2,31} = 2.872$, $p > 0.05$, Table 2), and thus grouped for analysis. Doses of 1.48 mg, 3.0 mg, and 5.0 mg were also tested. Differences in survival time were not significant at any dosage, however ($F_{0.05(1)4,63} = 0.544$, $p > 0.05$, Tables 4 and 5).

Two experiments were carried out on the extract (1.0 mg) of *J. mollissima*, the results of which were statistically homogeneous ($t_{0.05(2)10} = 0.991$, $p > 0.05$, Table 6) and thus grouped for comparison with the control group. As for the previous species, no significant difference was found in survival time ($t_{0.05(1)19} = 0.631$, $p > 0.05$, Table 6).

Finally, five experiments were conducted on the extract (1.0 mg) of *J. elliptica*, with doses of 0.74-10.0 mg, which were also homogeneous overall ($F_{0.05(1)4,24} = 1.972$, $p > 0.05$, Table 2). Once again, differences between the experimental and control

samples were not significant ($F_{0.05(1)4,39} = 0.91$, $p > 0.05$, Tables 7 and 8), contradicting the possible effect of the extract of this plant on the lethality of viper venom.

DISCUSSION

Plants known as “batata-do-teiú” – which means the “potato of the teiú lizard” (*Tupinambis* sp., family Teiidae) – are identified as containing an antivenin based on a legendary lizard which ate the root of the plant and thus became immune to the venom of vipers and rattlesnakes, *Bothrops* and *Crotalus* (Braga, 1960; Mors, 1991). A number of different plant species are known by this name (Martz, 1992), including *Apodanthera villosa*, the extract of which (1.0 mg) was shown in the present study to have a significant retardant effect on the lethality time of the animals inoculated with viper venom. No information is available on the chemical composition of this plant, nor experimental data on its antivenin properties.

The extracts of *Apodanthera glaziovii*, *Jatropha molissima* and *J. elliptica* not present effects of neutralization on the lethality of *Bothrops jararaca* venom.

There are numerous reports of plants that act on the venom of snakes, in particular the rattlesnakes, genus *Crotalus* (Mors *et al.*, 1989; Pithayanukul *et al.*, 2004). Nakagawa *et al.* (1982) pioneering experimental study of plants that affect the lethality of viper venom isolated substances denominated cabenegrin I and II from a plant identified only as “cabeça-de-negro”, supposedly Amazonian in origin. Cabenegrins are pterocarpanes of the isoflavonoid group, which are produced by plants as a defense against infection by virus or fungi (Costa, 2000). A plant with this name was also mentioned by Borges *et al.* (1996) in their analysis of the product “Específico Pessoa”, and concluded that it was ineffective for the treatment of the venom of *Bothrops atrox*. This word, “específico” is used generically for remedies in

homeopathy or to treat specific problems, and this product “Pessoa” – which is produced from a plant called “cabeça-de-negro”, according to its label – is a popular antivenin in northern and northeastern Brasil.

Cabenegrins were mentioned in a second study of plants with antivenin properties (Reyes-Chilpa *et al.*, 1994). The extract of the legume *Brongniartia podalyrioides* reduced mortality in mice inoculated with the venom of *Bothrops atrox*, in a similar form to that observed in the case of *A. villosa* in the present study. The active compound isolated from this species was the pterocarpan (-)-edunol, which related structurally to (-)-cabenegrins A-I and A-II.

Few reports are available on the composition of vegetable compounds that reduce the lethality of viper venom. One of the early studies of plants that affect viper venom records the action of the extract of *Mandevilla velutina* (Apocynaceae) on bradycinin, an endogenous polypeptide released into the blood plasma by the enzymatic action of viper venom (Calixto *et al.*, 1985). Pereira *et al.* (1994) identify a number of plants and their components as having an effect on the venom of *Bothrops jararaca*: i) triterpenes and steroids of *Periandra mediterranea* (Vell.) Taub (Fabaceae) and *Apuleia leiocarpa* (Vogt) Macbr (Caesalpinaceae); ii) derivatives of the cafeic acid of *Vernonia condensata* Baker (Compositae) and *Cynara scolymus* L. (Asteraceae); iii) coumarins of *Mikania glomerata* Spreng (Asteraceae) and *Dorstenia brasiliensis* Lam. (Moraceae); iv) flavonoids of *Phyllanthus klotzschianus* M. Arg (Euphorbiaceae), *Citrus sinensis* (L.) Osbeck (Rutaceae), *Apuleia leiocarpa* and *Derris sericea* (H.B.K.) Ducke (Fabaceae); v) lignoflavonoids of *Silybum marianum* Gaertn. (Compositae); vi) cumestans of *Eclipta prostrata*, vii) saponins of *Bredemeyera floribunda* Willd (Polygalaceae).

Mors *et al.* (2000) refer to a variety of substances isolated from plants which interfere in the lethal properties of the venom of *B. jararaca* in mice, such as corticosteroids, triterpenes, phenolic

compounds, hydroxybenzoic acids, chlorogenic acids, curcuminoids, coumarins, flavonoids, pterocarpanes, aristolochic acids, tannins and polysaccharides. All these substances belong to the same class of "secondary metabolites", which are capable of interacting with venom receptor and enzymes.

More recently, Ticli *et al.* (2005) analyzed the possible neutralization effects of the methanolic extract of *Cordia verbenacea* (Boraginaceae) on the venom of *Bothrops jararacussu* and its principal phospholipases A₂ (Bothropstoxins I and II). They isolated the plant's active ingredient, identified as rosmarinic acid, and concluded that the pure compound potentializes the action of viper-rattlesnake antivenin by neutralizing the venom's lethal effects.

REFERENCES

- Amuá, S.F., S. Marcussi, M. Urzêda, L.B. Silveira, V.C. Fernandes, M.F.B. Coelho, A.M.S. Pereira, S.C. França & A.M. Soares, 2003. Atividade antiofídica de extratos de *Jatropha elliptica* (Pohl.) Muell. Arg. **VI Jornada Paulista de Plantas Medicinais**, São Pedro, São Paulo.
- Borges, C.C., 1996. Eficácia da espécie vegetal *Peltodon radicans* Phl. (Labiatae, Lamiaceae) na neutralização da atividade edematogênica e na ineficácia do extrato vegetal Específico Pessoa na neutralização das principais atividades do veneno de *B. atrox*. **Revista da Universidade Federal do Amazonas, Série Ciências Biológicas** 1: 97-113.
- Braga, R., 1960. Plantas do nordeste, especialmente do Ceará. 2ª edição. **Imprensa Oficial**, Fortaleza 540p.
- Calixto, J.B., M. Nicolau & R.A. Yunes, 1985. The selective antagonism of bradykinin action on rat isolated uterus by crude *Mandevilla velutina* extract. **Journal of Brazilian Pharmacology** 85(4):729-731.
- Costa, R.R.P. 2000. Sofrol e eugenol: estudo da reatividade química e uso em síntese de produtos naturais biologicamente ativos e seus derivados. **Química Nova** 23(3):357-369.
- Fisher, R.A., 1949. **The design of experiments**. Fifth edition. Oliver and Boyd, Edinburg 242p.
- Fisher, R.A. & F. Yates, 1953. **Statistical tables for biological agricultural and medical research**. Fourth edition. Oliver and Boyd, Edinburgh 126p.
- Furtado, M.F.D., G.M.D.D. Colleto & W.D. Silva, 1991. Controle de qualidade dos venenos animais e dos correspondentes antivenenos. I. Padronização dos métodos de ensaios das atividades bioquímicas e farmacológicas dos venenos de algumas espécies do gênero *Bothrops* e *Crotalus* usando amostras secas a temperatura ambiente ou liofilizadas. **Memórias do Instituto Butantan** 53(2):149-159.
- Gutiérrez, J.M. 2002. Compreendendo los venenos de serpientes: 50 años de investigaciones em América Latina. **Revista de Biología Tropical** 50(2):377-394.
- Martz, W., 1992. Plants with a reputation against snakebite. **Toxicon** 30 (10):1131-1142.
- Mors, W.B., 1991. Plant active against snake bite, pp. 353-373. In: **Economic and medicinal plant research** (Wagner, H., H. Kihino & N.R. Farnsworth, Eds.). Vol. 5. Academic Press, New York.
- Mors, W.B., M.C. Nascimento, J.P. Parente, M.H. Silva, P.A. Melo & G. Suarez-Kurtz, 1989. Neutralization of lethal and mitotoxic activities of south american rattlesnake venom by extracts and constituents of the plant *Eclipta prostrata* (Asteraceae). **Toxicon** 27(9):1003-1009.
- Mors, W.B., Nascimento, M.C., Pereira, B.M.R., Pereira, N.A., 2000. Plant natural products active against snake bite – the molecular approach. **Phytochemistry** 55:627-642.
- Nakagawa, M., K. Nakanishi, L.L. Darko & J.A. Vick, 1982. Structures of cabenegrins A-I and A-II, potent anti-snake venoms. **Tetrahedron Letters** 23 (38):3855-3858.
- Otero, R., R. J. Fonnegra & S.L. Jiménez, 2000. **Plantas utilizadas contra mordeduras de serpientes en Antioquia y Chocó, Colombia**. Universidad de Antioquia, Medellín 402p.
- Pereira, N.A., B.M.R. Pereira, M.C. Nascimento, J.P. Parente & W.B. Mors, 1994. Pharmacological screening of plants recommended by folk medicine as snake venom antidotes; IV. Protection against jararaca venom by isolated constituents. **Planta Medica**, Stuttgart 60: 99-100.
- Pithayanukul, P., S. Laovachirasuan, R. Bovovada, N. Pakmanee & R. Suttisri, 2004. Anti-venom potential of butanolic extract of *Eclipta prostrata* against Malayan pit viper venom. **Journal of Ethnopharmacology** 90:347-352.
- Reyes-Chilpa, R., F. Gómez-Garibay, L. Quijano, G.A. Magos-Guerrero & T. Rios, 1994. Preliminary results on the protective effect of (-)-edunol, a pterocarpin from *Brongniartia podalyrioides* (Leguminosae), against *Bothrops atrox* venom in mice. **Journal of Ethnopharmacology** 42: 199-203.
- Silva, G.L., M.I.L. Machado, F.J.A. Matos & R. Braz-Filho, 1999. A new isolated from *Harpalyce brasiliana*. **Journal of the Brazilian Chemical Society** 10(6):438-442.
- Ticli, F.K., L.I.S. Hage, R.S. Cambráia, P.S. Pereira, A.I. Magro, M.R.M. Fontes, R.G. Stábili, J.R. Giglio, S.C. França, A.M. Soares & S.V. Sampaio, 2005. Rosmarinic acid, a new snake venom phospholipase A₂ inhibitor from *Cordia verbenacea* (Boraginaceae):

- antiserum action potentiation and molecular interaction. **Toxicon** 46:318-327.
- Vanzolini, P.E., 1993. **Métodos estatísticos elementares em sistemática zoológica**. Hucitec, São Paulo 130p.
- Vilar, J.C., C.M. Carvalho & M.F.D. Furtado, 2004. Epidemiologia dos acidentes ofídicos em Sergipe (1999-2002). **Biologia Geral e Experimental** 4(2):3-13.
- Vilar, J.C., C.M. Carvalho & M.F.D. Furtado, 2005. Ofidismo e plantas utilizadas como antiofídicas. **Biologia Geral e Experimental** 6(1):3-36.
- Zar, J.H., 1996. **Biostatistical Analysis**. Third edition. Prentice Hall, New Jersey 662p.

Aceito: 12.vi.2007

Table 1. Regression of the probitic value (y) on the logarithm of the concentration (x) of the 50% lethal dose (LD₅₀), control (μg) and extracts (mg).

	y = a+bx	ly	LD ₅₀
Control	- 7.99 + 8.28x	34.43±40.45	37.09
<i>Apodanthera villosa</i>	10.797+3.358x	0.015±0.023	0.018
<i>Apodanthera glaziovii</i>	12.722+3.596x	0.0059±0.0089	0.007
<i>Jatropha mollissima</i>	9.910+1.831x	0.0013±0.003	0.002
<i>Jatropha elliptica</i>	15.691+6.193x	0.013±0.023	0.018

b, regression coefficient
 a, constant of regression

ly, confidence interval of y
 LD₅₀ for y = 5

Table 2. ANOVA of survival (hours) of mice and homogeneity of control and experimental samples.

Control/Extract	Source of variation	Degrees of freedom	Sum of squares	Mean square	F
Control	Groups	5	5.721	1.144	1.436ns
	Error	27	21.52	0.7971	
<i>Apodanthera villosa</i>	Groups	3	529.95	176.65	0.857ns
	Error	42	8650.7	205.97	
<i>Apodanthera glaziovii</i>	Groups	2	311.20	155.60	2.872ns
	Error	31	1679.6	54.182	
<i>Jatropha elliptica</i>	Groups	4	2.433	0.6082	1.972ns
	Error	24	7.401	0.3084	

Table 3. Statistics of the frequency distribution of mouse survival (hours), and comparison of the means of the control and sample treated with the extract of *Apodanthera villosa*.

Group	Dose of venom (μg)	Dose of extract (mg)	N	A	m	s	CV	Im	t
Control	74.2	-	23	2.26-5.93	3.203±0.182	0.875	27.31	2.825-3.581	2.084 *
Extract	74.2	1.0	46	1.56-48.0	7.609±2.106	14.283	187.7	3.364-11.854	

N, sample size

m, mean±standard error

* significant at 5% level

A, amplitude

CV, coefficient of variation

Im, confidence interval of the mean

s, standard deviation

Table 4. Statistics of the frequency distribution of mouse survival (hours), and comparison of the means of the control and sample treated with extracts of *Apodanthera glaziovii*.

Group	Dose of venom (μg)	Dose of extract (mg)	N	A	m	s	CV	Im
Control	74.2	-	16	2.26-5.93	3.614±0.284	1.138	31.48	3.00-4.221
Extract (1)	74.2	1.00	34	2.05-48.0	4.201±1.332	7.767	184.8	1.489-6.912
Extract (2)	74.2	1.48	6	2.16	2.16±0.0	0.0	0.0	2.16
Extract (3)	74.2	3.00	6	1.35-3.45	2.027±0.3	0.737	36.35	1.253-2.8
Extract (4)	74.2	5.00	6	1.06-1.56	1.297±0.08	0.218	16.80	1.068-1.526

N, sample size

m, mean±standard error

() Experiments 1-4

A, amplitude

CV, coefficient of variation

Im, confidence interval of the mean

s, standard deviation

Table 5. ANOVA of survival (hours) of mice and homogeneity of control and experimental samples of *Apodanthera glaziovii*.

Source of variation	Degree of freedom	Sum of squares	Mean squared	F
Group	4	69.532	17.383	0.544 ns
Error	63	2013.2	31.956	

ns, not significant

Table 6. Statistics of the frequency distribution of mouse survival (hours), control and extract of *Jatropha mollissima*, comparison of the means of the experiments 1 and 2 of the extract and between the control and extract.

Group	Dose of venom (μ g)	Dose of extract (mg)	N	A	m	s	CV	Im	t
Extract 1	74.2	-	6	2.35-48.0	10.01 \pm 7.597	18.608	185.7	-9.51-29.54	0.991ns
Extract 2	74.2	1.0	6	1.55-3.33	2.485 \pm 0.254	0.623	25.07	1.83-3.14	
Control	74.2	-	10	2.70-5.13	3.758 \pm 0.268	0.849	22.59	3.151-4.365	0.63 ns
Extract	74.2	1.0	11	1.55-48.0	6.517 \pm 4.15	13.763	211.1	-2.728-15.763	

N, sample size
m, mean \pm standard error
ns, not significant

A, amplitude
s, standard deviation
CV, coefficient of variation
Im, confidence interval of the mean

Table 7. Statistics of the frequency distribution of mouse survival (hours), comparison of the means of the control and sample treated with extracts of *Jatropha elliptica*.

Group	Dose of venom (μ g)	Dose of extract (mg)	N	A	m	s	CV	Im
Controle	74.2	-	11	2.66-4.93	3.325 \pm 0.202	0.670	20.15	2.874-3.775
Extract (1)	74.2	0.74	6	2.25-3.26	2.945 \pm 0.144	0.353	11.98	2.574-3.316
Extract (2)	74.2	1.00	6	2.00-3.53	2.842 \pm 0.207	0.508	17.87	2.308-3.375
Extract (3)	74.2	1.48	6	1.63-3.38	2.73 \pm 0.247	0.605	22.16	2.094-3.366
Extract (4)	74.2	5.00	8	1.48-48.0	7.841 \pm 5.74	16.236	207.0	-5.734-1.417
Extract (5)	74.2	10.00	8	1.4-2.68	2.083 \pm 0.196	0.555	26.64	1.618-2.547

N, sample size
m, mean \pm standard error
() Experiments 1-5

A, amplitude
s, standard deviation
CV, coefficient of variation
Im, confidence interval of the mean

Table 8. ANOVA of the survival time (hours) of mouse, control and extract of *Jatropha elliptica*.

Source of variation	Degree of freedom	Sum of squares	Mean squared	F
Grupos	5	173.16	34.632	0.727 ns
Erro	39	1855.6	47.580	

ns, not significant